

PACKING A CAKE INTO A BOX

M. SKOPENKOV

ABSTRACT. Given a cake in form of a triangle and a box that fits the mirror image of the cake, how to cut the cake into a minimal number of pieces so that it can be put into the box? The cake has an icing, so that we are not allowed to put it into the box upside down. V.G. Boltyansky asked this question in 1977 and showed that three pieces always suffice. In this paper we provide examples of cakes that cannot be cut into two pieces to put into the box. This shows that three is the answer to V.G. Boltyansky's question. Also we give examples of cakes which can be cut into two pieces.

1. EXAMPLES

Given a cake in form of a triangle and a box that fits the mirror image of the cake, how to cut the cake into a minimal number of pieces so that it can be put into the box? The cake has an icing, so that we are not allowed to put it into the box upside down, see Figure 1.



FIGURE 1. A cake and a box

V.G. Boltyansky asked this question in his book on Hilbert's third problem [1, §9]. Let us give an accurate statement, see Figure 2. A *cake* $\triangle ABC$ is a nonisosceles triangle in the plane with vertices A, B, C and angles α, β, γ . A *box* $\triangle A'B'C'$ is a triangle obtained from the cake by a reflection with respect to a line. A cutting of the cake is *nice* if the pieces are polygons which can be put into the box all together by means of appropriate rotations and translations. The problem we study is to cut the cake nicely into a minimal number of pieces.




FIGURE 2. A cake and a box again

V.G. Boltyansky gives the following example:

Example 1. [1, §9] Any cake can be nicely cut into **3** pieces.



FIGURE 3. A nice cutting into 3 pieces

Proof. Indeed, cut the cake along the perpendiculars from the incenter to the sides, see Figure 3. Rotating the pieces around the incenter pack the cake into the box. $\square$

This is all concerning the question we find in the book. However, V.V. Prasolov indicated that some cakes (see Example 2(a)–(d)) can be nicely cut into 2 pieces.

Example 2. Cakes of indicated shapes (see Figure 4) can be nicely cut into 2 pieces:

- (a) $\alpha = 90^\circ$; (b) $\alpha = 3\beta$; (c) $\alpha = 2\beta < 90^\circ$; (d) $\alpha = 2\beta > 90^\circ$;
 (e) $\alpha = 30^\circ$, $\beta = 20^\circ$, $\gamma = 130^\circ$; (f) $\alpha = \frac{n+1}{n}\beta$, where n is an integer.




FIGURE 4. Cakes which can be nicely cut into 2 pieces

We are not going to do the reader out of a pleasure to cut the cakes in Examples (a)–(d) nicely himself (herself) by means of straight cuts. Answers are given in §5.

Let us present two ways of cutting the cake in Example (e), which is probably new.

1. A wheel-shaped nice cut. A cake with $\alpha : \beta = 3 : 2$ can be nicely cut as follows, see Figure 5. Take a broken line $AKLMNB$ with 5 equal sides and equal angles $180^\circ - \alpha + \beta$ between each pair of consecutive sides. Clearly, the hexagon $AKLMNB$ is mirror-symmetric. Since the sum of the angles of the hexagon is 720° it follows that $\angle BAK = \angle ABN = \beta$. Thus $N \in BC$ (assume that both points N and C are above the line AB). The cut $AKLMN$ is nice. Indeed, $\angle CNM = \alpha - \beta = \angle CAK$. Thus the hexagon $AKLMNC$ is also mirror-symmetric and $AC = NC$. Rotating the "wheel" $AKLMNB$ one packs the cake into the box. $\square$

2. A gear-shaped nice cut. A cake with $\alpha : \beta = 3 : 2$ can be also nicely cut as follows, see Figure 6. Take a saw-shaped broken line $AKLMNB$ with 5 equal sides and angles $\angle AKL = \angle LMN = 180^\circ - \beta$, $\angle KLM = \angle MNB = 180^\circ - \alpha$. Then $K \in AB$ and $N \in BC$. The cut $KLMN$ is nice: rotating the "gear" $AKLMNB$ one packs the cake into the box. $\square$

There is even one more way to cut the cake of Example (e). The reader may wish to find it himself (herself) or see the answer in §5.

Similar wheel- and gear-shaped cuts nicely cut the cake of Example (f), see Figures 7 and 8.

Conversely, if the cut in Figure 5 is nice then $\alpha : \beta = 3 : 2$. Indeed, assume that the piece $AKLMNB$ is put into the box by a rotation R such that $R(B) = N = A'$ and $C = C'$. Then $R(N) = M$, $R(M) = L$, $R(L) = K$, $R(K) = A$. A computation shows that the hexagon $AKLMNB$ has 2 angles of size β and 4 angles of size $180^\circ - \alpha + \beta$. Since their sum is 720° it follows that $2\alpha - 3\beta = 0$. This is a particular case of the following theorem (announced in [2]):

Theorem 3. *If a cake can be nicely cut into 2 pieces then $k\alpha + l\beta + m\gamma = 0$ for some integers k , l and m , not vanishing simultaneously.*

Thus *almost all* cakes cannot be nicely cut into 2 pieces (e.g., a cake with $\alpha = \sqrt{2}^\circ$, $\beta = \sqrt{3}^\circ$, $\gamma = 180 - \sqrt{2} - \sqrt{3}^\circ$).

One can try to prove Theorem 3 analogously to the above proof that $2\alpha - 3\beta = 0$ in Figure 5. But this attempt leads to a huge exhaustion. In this paper we give a short elementary proof of the theorem.




FIGURE 5. A wheel-shaped nice cut




FIGURE 6. A gear-shaped nice cut




FIGURE 7. A wheel-shaped nice cut again




FIGURE 8. A gear-shaped nice cut again

2. INVARIANTS

Let us discuss historical background of Theorem 3. Our history begins with

Theorem (Bolyai–Gerwien, 1832). *Suppose that two planar polygons have the same area. Then the first polygon can be cut into finitely many polygonal pieces which can be reassembled as the second polygon (possibly turning over the pieces).*

V.G. Boltyansky comes to his question while proving that the Bolyai–Gerwien theorem remains true if turning over the pieces is not permitted. Let us sketch the proof. Assume that a piece P is turned over during the assembling. Triangulate the piece. Cut each of the obtained triangles into 3 subpieces as in Example 1. These subpieces can be reassembled as the mirror image of P . Thus one gets rid of turnings.

H. Hadwiger and P. Glur go even further. They prove that Bolyai–Gerwien theorem remains true if only parallel translations and central symmetries of the pieces are permitted [1, §9]. The latter result is optimal: this collection of isometries is in some sense minimal [1, §10]. The proof of optimality uses an *additive invariant*, which is also the main tool of the paper.

Definition (of the additive invariant). [1, §10] Let P be a polygon in the plane, see Figure 9. Orient all the sides of the polygon P counterclockwise. Let f be a *directed line*, i.e., a line with a direction on it marked by an arrow. The *additive invariant* $J_f(P)$ is an algebraic sum of all the sides of the polygon P parallel to f . In this sum the sides having the *same* direction as f (sides AB , CD and FG in Figure 9), are taken with positive sign, and the ones having the *opposite* direction (side JK in Figure 9) are taken with negative sign. If the polygon P has no sides parallel to f then set $J_f(P) = 0$.




FIGURE 9. Definition of the additive invariant: $J_f(P) = AB + CD + FG - JK$

Let us establish some basic properties of the additive invariant.

Claim 4. [1, §10] (a) *Suppose that a polygon P is cut into several polygons $P_1, \dots, P_n$. Then $J_f(P) = J_f(P_1) + \dots + J_f(P_n)$.*

(b) *Suppose that a cake ∇ is cut into several polygonal pieces which are put into the box ∇ all together by means of parallel translations. Then $J_f(\nabla) = J_f(\nabla)$.*

Proof. [1, §10] (a) Mark all the vertices of the polygons $P, P_1, \dots, P_n$. Marked points split the sides of the polygons into smaller segments called *edges*, see Figure 10. Each side is the sum of the edges, into which the side splits. Thus the algebraic sum of the *sides* can be replaced by the algebraic sum of the *edges* in the definition of the additive invariant.

Consider the sum $J_f(P_1) + \dots + J_f(P_n)$. It equals to the algebraic sum of all the edges parallel to f of all the polygons $P_1, P_2, \dots, P_n$.

Take an edge inside the polygon P (possibly excluding the endpoints). Then there are two polygons P_i and P_j bordering upon the edge from opposite sides. The contributions of the edge to $J_f(P_i)$ and $J_f(P_j)$ have opposite signs and hence cancel in the sum $J_f(P_1) + \dots + J_f(P_n)$.

Take an edge in the boundary of the polygon P . Then there is a unique polygon P_i bordering upon the edge. The polygon P_i borders upon the edge from the same side as the polygon P . Thus the contributions of the edge to $J_f(P_i)$ and $J_f(P)$ are the same. Hence $J_f(P_1) + \dots + J_f(P_n) = J_f(P)$.




FIGURE 10. Additivity of the invariant: $J_f(P) = J_f(P_1) + J_f(P_2) + J_f(P_3)$

(b) Suppose that the cake ∇ is cut into polygonal pieces $P_1, \dots, P_n$, the box ∇ is cut into polygonal pieces $P'_1, \dots, P'_n$, and the piece P'_i is a parallel translation of the piece P_i for each $i = 1, \dots, n$. Clearly, $J_f(P_i) = J_f(P'_i)$. Thus by assertion (a) we have

$$J_f(\nabla) = J_f(P_1) + \dots + J_f(P_n) = J_f(P'_1) + \dots + J_f(P'_n) = J_f(\nabla).$$

□

To explain our idea let us prove an assertion close to Theorem 3. This assertion also shows that the Bolyai–Gerwien theorem is not necessarily true if only parallel translations of the pieces are permitted.

Claim 5. *It is impossible to cut a cake into polygonal pieces and put the pieces into a box all together by means of only parallel translations.*

Proof. Assume that the cake is cut as required. Then $J_{\mathbf{AB}}(\nabla) = AB$. By Claim 4(b) $J_{\mathbf{AB}}(\nabla) = J_{\mathbf{AB}}(\nabla)$. Thus $J_{\mathbf{AB}}(\nabla) \neq 0$. Hence the box has a side parallel to AB . Analogously the box has 2 sides parallel to BC and CA . Thus the box is either a parallel translation of the cake or central-symmetric to the cake. Since the box is a mirror image of the cake it follows that the cake is mirror-symmetric, i.e., isosceles. This contradiction proves the claim. □

For the proof of Theorem 3 we need to generalize the invariant J_f to make it invariant under a rotation. Denote by $\mathbf{XY}$ the directed line passing through points $X \neq Y$ and directed from X to Y .

Definition (of the generalized additive invariant). Let $f(\mathbf{XY})$ be a function on the set of all directed lines $\mathbf{XY}$ in the plane satisfying the property: $f(\mathbf{XY}) = -f(\mathbf{YX})$. Let $P = A_1A_2 \dots A_n$ be a polygon whose vertices are enumerated counterclockwise. Set

$$J_f(P) = f(\mathbf{A_1A_2})A_1A_2 + f(\mathbf{A_2A_3})A_2A_3 + \dots + f(\mathbf{A_nA_1})A_nA_1.$$

We say that a rotation R preserves the function f if for each directed line $\mathbf{XY}$ we have $f(R(\mathbf{XY})) = f(\mathbf{XY})$. If R preserves f then R preserves the invariant J_f , i.e., $J_f(R(P)) = J_f(P)$ for any polygon P . The following claim is proved analogously to Claim 4.

Claim 6. *Suppose that a cake ∇ is cut into several polygons and the pieces are put into a box ∇ all together by means of rotations preserving the function f . Then $J_f(\nabla) = J_f(\nabla)$.*

3. PROOF OF THE THEOREM

We are ready to prove Theorem 3.

Let the cake ∇ be nicely cut into 2 pieces. We may assume that while packing into the box $\triangledown$ one piece remains fixed and the other piece is moved by a rotation or parallel translation R . By Claim 5 it follows that R is not a parallel translation. Thus R is a rotation through an angle ϕ around a point O , see Figure 11.




FIGURE 11. The rotation R

We say that an angle ψ is a *multiple* of ϕ , if $\psi = k\phi + l \cdot 180^\circ$ for some integers k and l . Notation: $\phi \mid \psi$. An angle ψ is *rational* if $\psi = k \cdot 180^\circ / l$ for some integers k and l .

Our aim is to show that one of the following conditions holds:

- each of the angles $2\alpha, 2\beta, 2\gamma$ is a multiple of ϕ (as in Example 2(e)); or
- one of the angles $2\alpha, 2\beta, 2\gamma$ is a multiple of ϕ and ϕ is rational (as in Example 2(a)).

The *oriented angle* $\angle(KL, MN)$ between (undirected) lines KL and MN is an angle of a counterclockwise rotation taking the line KL to MN . An oriented angle is defined up to a multiple of 180° .

Claim 7. *If either ϕ is irrational or $O \notin \mathbf{AB}$ then $\phi \mid \angle(AB, A'B')$.*

Proof. Let us construct an invariant J_f preserved by the rotation R . Denote by $R^1, R^2, R^3, \dots$ the iterations of the rotation R , by $R^{-1}, R^{-2}, R^{-3}, \dots$ the iterations of the inverse rotation R^{-1} , and by R^0 the identity map. Define a function f on the set of all directed lines by the formula

$$f(\mathbf{XY}) = \begin{cases} 1, & \text{if } \mathbf{XY} = R^k(\mathbf{AB}) \text{ for some integer } k; \\ -1, & \text{if } \mathbf{XY} = R^k(\mathbf{BA}) \text{ for some integer } k; \\ 0, & \text{otherwise.} \end{cases}$$

For instance, in Figure 11 $f(\mathbf{AB}) = 1$, $f(\mathbf{A'B'}) = -1$, $f(\mathbf{AC}) = 0$.

Since either ϕ is irrational or $O \notin \mathbf{AB}$ it follows that $R^k(\mathbf{AB}) \neq R^l(\mathbf{BA})$ for all integers k, l . Thus the function f is well-defined. Clearly, the rotation R preserves the function f . Notice that the condition $f(\mathbf{XY}) \neq 0$ implies $\phi \mid \angle(AB, XY)$.

Consider the invariants $J_f(\nabla)$ and $J_f(\triangledown)$. Both of them are linear combinations of the sides AB, BC, CA with coefficients $0, \pm 1$. For instance, in Figure 11 $J_f(\nabla) = J_f(\triangledown) = AB + BC$.

Let us prove that at least one of the sides AB, BC, CA has nonzero contribution to both $J_f(\nabla)$ and $J_f(\triangledown)$. Indeed, otherwise the difference $J_f(\nabla) - J_f(\triangledown)$ is also a linear combination of these sides with coefficients $0, \pm 1$. Such a linear combination is nonzero by the triangle inequality and the assumption that the cake is not isosceles. On the other hand, by Claim 6 we have $J_f(\nabla) - J_f(\triangledown) = 0$, a contradiction.

So a side of the cake has nonzero contribution to both $J_f(\nabla)$ and $J_f(\triangledown)$.

In case when the side is AB we get $f(\mathbf{A'B'}) \neq 0$. Thus $\phi \mid \angle(AB, A'B')$.

In case when the side is BC we get $f(\mathbf{BC}) \neq 0$ and $f(\mathbf{B'C'}) \neq 0$. Thus $\phi \mid \angle(AB, BC)$ and $\phi \mid \angle(AB, B'C')$. Then

$$\angle(AB, A'B') = \angle(AB, B'C') + \angle(B'C', A'B') = \angle(AB, B'C') + \angle(AB, BC)$$

is a multiple of ϕ .

Case when the side is AC is analogous. □

Claim 8. *If either ϕ is irrational or $O \notin \mathbf{AB}, \mathbf{AC}$ then $\phi \mid 2\alpha$.*

Proof. By Claim 7 we have $\phi \mid \angle(AB, A'B')$ and $\phi \mid \angle(AC, A'C')$. Thus

$$2\alpha = \angle(AB, AC) + \angle(A'C', A'B') = \angle(AB, A'B') - \angle(AC, A'C')$$

is a multiple of ϕ . □

Claim 9. *At least two lines containing the sides of the cake do not pass through the point O .*

Proof. Assume the converse. Then O is a vertex of the cake. Assume without loss of generality that $O = A$ and $AB > AC$.

Let us show that $A' = A$. Indeed, consider all the angles with vertex at A belonging to both pieces. The sum of the angles is α and is preserved by the rotation R . Thus the box has angle of size α at the point A . That is, $A' = A$.

Let us show that either $B' = B$ or $B' = R(B)$. Indeed, the point B of the cake is the most distant from A . Thus no other points of the cake border upon B after the rearrangement of the pieces. Thus the box has angle of size β at the point B or $R(B)$, depending on whether the piece containing B is fixed or mobile. That is, either $B' = B$ or $B' = R(B)$.

In case $A' = A$ and $B' = B$ we get $\nabla \cap \triangleright = A'B'$. Thus the fixed piece is empty.

In case $A' = A$ and $B' = R(B)$ we get $R(\nabla) \cap \triangleright = A'B'$. Thus the mobile piece is empty.

This contradiction proves the claim. □

Now the proof of the theorem is concluded by consideration of the following 2 cases.

Case 1: ϕ is irrational. By Claim 8 it follows that $\phi \mid 2\alpha, 2\beta, 2\gamma$. Then $k\alpha + l\beta + m\gamma = 0$ for some integers k, l and m , not vanishing simultaneously.

Case 2: ϕ is rational. By Claim 9 we may assume without loss of generality that $O \notin \mathbf{AB}, \mathbf{AC}$. Then by Claim 8 $\phi \mid 2\alpha$. Thus $\alpha = k \cdot 180^\circ / l$ for some integers k and l . So $0 = k \cdot 180^\circ - l\alpha = (k - l)\alpha + k\beta + k\gamma$.

4. AN OPEN PROBLEM

Let us state an open problem, cf. [3]. We say that two similar nonisosceles triangles in the plane are *oriented oppositely* if one of them includes angles α, β, γ clockwise, and another one counterclockwise, see Figure 2. For instance, the height from the right angle cuts a right triangle into two triangles similar to it but oriented oppositely.

Problem 10. Does there exist a nonright and nonisosceles triangle which can be cut into triangles similar to it but oriented oppositely?

Let us announce a partial result due to M.V. Prasolov and the author. The proof is based on a similar invariant as above.

Theorem 11. *Suppose that a triangle with angles α, β and γ can be dissected into triangles similar to it but oriented oppositely. Then $k\alpha + l\beta + m\gamma = 0$ for some integers k, l and m , not vanishing simultaneously.*

5. ANSWERS

Proof of Example 2(a)–(d). A useful observation is: if the pieces are mirror-symmetric then the cutting is automatically nice. This allows not to take care about rearrangement of the pieces.

(a) Cut the cake along the median from the vertex of the right angle, see Figure 12(a).

(b) Cut the cake along the line separating an angle of size β from the angle α , see Figure 12(b).

(c) Cut the cake along the line separating an angle of size β from the angle γ , see Figure 12(c).

(d) Cut the cake along the line symmetric to the side opposite to the angle γ with respect to the bisector of the angle γ , see Figure 12(d). $\square$




FIGURE 12. Nice cuttings of cakes from Figure 4




FIGURE 13. A scissors-shaped nice cut

Third proof of Example 2(e): **3. A scissors-shaped nice cut.** Take a broken line $BCKLMB$ with $CK = KL = LM$ and with equal angles 130° between consecutive edges, see Figure 13. Then $K \in AC$, $M \in AB$ and the cut KLM is nice. Rotating the "scissors" $AKLM$ and $BCKLMB$ we pack the cake into the box. $\square$

6. ACKNOWLEDGEMENTS

The author is grateful to B.R. Frenkin, A.A. Glazyrin, I.V. Izvestiev and M.V. Prasolov for useful discussions. The author is also grateful to his wife Anastasia for some figures and cakes. The author was supported in part by Moebius Contest Foundation for Young Scientists and the Euler Foundation.

REFERENCES

- [1] V.G. Boltianskii, *Hilbert's third problem*, Transl. by R. Silverman, V. H. Winston & Sons, Washington D.C., 1978.
- [2] M. Prasolov, M. Skopenkov and B. Frenkin, Invariants of polygons, *XIX Summer conference of the International mathematical tournament of towns*, 2007, <http://turgor.ru/lktg/2007/1/1-1en.pdf>.
- [3] A. Zak, Dissection of a triangle into similar triangles, *Discr. Comp. Geom.* **34:2** (2005), pp. 295–312.

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS OF THE RUSSIAN ACADEMY OF SCIENCES, BOLSHOY KARETNY PER.
19, BLD. 1, MOSCOW, 127994, RUSSIAN FEDERATION, AND KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY,
4700 THUWAL, 23955-6900, KINGDOM OF SAUDI ARABIA
E-mail address: `skopenkov@rambler.ru`